

Instructions After a Tooth Extraction

DO NOT DISTURB THE AREA: For the next few days, and especially the first 24 hours, it is very important to allow your body to form a good clot and start the natural healing process. Swishing, sucking through a straw, and smoking can all dislodge the clot. Keep anything sharp from entering the wound (crunchy food, toothpicks, eating utensils). Be sure to chew on the opposite side for 24 hours.

BLEEDING: When you leave the office, you will be biting on a gauze pad to control bleeding. Keep slight pressure on this gauze for at least 15-20 minutes. Don't change it during this time; it needs to remain undisturbed while a clot forms in the extraction socket. You may bite on another gauze or a tea bag for another 30 minutes if you feel it is still bleeding. Constant pressure on the area will stop the bleeding. Small amounts of blood in the saliva can make your saliva appear quite red. This is normal and may be noticed the rest of the day after the procedure.

PAIN: Some discomfort is normal after surgery. If prescription pain medication is prescribed, take it as instructed on the label. Don't exceed the dose on the label. Taking with food or milk will help reduce upset stomach. Avoid driving or operating heavy machinery when taking pain prescriptions. Do not drink alcohol while taking prescription pain medications. If you do not react well to the prescription, or if one was not prescribed take two Tylenol, Advil, or similar non-aspirin pain reliever every 3 to 4 hours until bedtime to maintain comfort. Take it before the anesthesia wears off. Do not mix pain medications together.

SMOKING: Smoking should be stopped following surgery. Healing and success of the surgery will be substantially reduced by the cigarette smoke chemicals in your body. Smokers are at greater risk of developing a painful dry socket.

NAUSEA: This is most often caused by taking pain medications on an empty stomach. Reduce nausea by preceding each pain pill with soft food, and taking the pill with a large glass of water.

SWELLING: Applying an ice bag to the face over the operated area will minimize swelling. Apply for 15 minutes, then remove for 15 minutes. Continue this for the first day as needed.

NUMBNESS: The local anesthetic will cause you to be numb for several hours after you leave the office. Be very careful not to bite, chew, pinch, or scratch the numb area.

BRUSHING: Do not brush your teeth for the first 8 hours after surgery. After this, you may brush your teeth gently, but avoid the area of surgery for 3 days.

RINSING: Avoid all rinsing or swishing for 24 hours after extraction. Rinsing can disturb the formation of a healing blood clot which is essential to proper healing. This could cause bleeding and risk of a dry socket. After 24 hours you may begin gentle rinsing with a saltwater solution (1/2 teaspoon salt + 1/2 teaspoon soda + 8 ounces warm water). Avoid commercial mouth rinses.

DIET: Eat soft foods for the first two days. Maintain a good, balanced diet. Return to normal regular meals as soon as you are able after the first two days. Drink plenty of water. Avoid alcohol for 48 hours.

ACTIVITY: After leaving the office, rest and avoid strenuous activities for the remainder of the day. Keeping blood pressure lower will reduce bleeding and aid in healing.

ANTIBIOTICS: If you were given an antibiotic prescription, take all of them as directed until they are gone. Women: some antibiotics can reduce the effectiveness of birth control pills. Use alternate birth control methods for two months.

FOLLOW-UP APPOINTMENTS: You may need to return to the office to have sutures removed, or just for a brief follow-up healing check.

Please call Eastern Family Dentistry if you have:

- uncontrollable pain
- excessive or severe bleeding
- marked fever
- excessive warm swelling occurring a few days after the procedure
- reactions to medications, especially rash, itching, or breathing problems



Phone: (575) 356-8514